Statistical mechanics of a gas at high temperature

Statistical mechanics of a gas at “high” (i.e. not extremely low) temperature

1. Obtain partition function for the translational motion of particles in a box
2. Hence obtain energy, heat capacity, **and entropy**
3. **The idea of k-space and density of states**
4. Bring in rotation and vibration

Statistical mechanics of a gas at “high” (i.e. not extremely low) temperature

1. → Obtain partition function for the translational motion of particles in a box
2. Hence obtain energy, heat capacity, **and entropy**
3. **The idea of k-space and density of states**
4. Bring in rotation and vibration

For a particle moving in a one-dimensional potential well, there is **one quantum state of motion** (energy eigenstate) **for each volume h in position-momentum space**.



(We will prove this shortly for the infinite square well, but it is true more generally.)

Statistical mechanics of a gas at “high” (i.e. not extremely low) temperature

1. Obtain partition function for the translational motion of particles in a box
2. →Hence obtain energy, heat capacity, **and entropy**
3. **The idea of k-space and density of states**
4. Bring in rotation and vibration



Entropy of neon as a function of temperature,
with prediction of Sackur-Tetrode (i.e. stat. mech.) equation.



Entropy of neon as a function of temperature, with prediction of Sackur-Tetrode equation.

Entropy prediction
WRONG at low T





At low T we have to take **indistinguishability** into account, which we have NOT done here.

Entropy of neon as a function of temperature, with prediction of Sackur-Tetrode equation.

Statistical mechanics of a gas at “high” (i.e. not extremely low) temperature

1. Obtain partition function for the translational motion of particles in a box
2. Hence obtain energy, heat capacity, **and entropy**
3. **→ The idea of k-space and density of states**
4. Bring in rotation and vibration

Gas in a box





Potential well with

energy
eigenstates
(in one dimension)

Gas in a box





Potential well with

energy
eigenstates
(in one dimension)



States in k space

Each dot is an allowed state at some energy

The quantity

$$\frac{1}{\Delta k_x \, \Delta k_y \, \Delta k_z} = \frac{V}{\pi^3}$$

is called the **density of states in k-space**.

The quantity

$$\frac{\mathrm{d}N}{\mathrm{d}k}$$

is also a **density of states**, but now a density with respect to the magnitude of k.

This is often written $g(k) \equiv \frac{\mathrm{d}N}{\mathrm{d}k}$ and so we have:

$$(\text{number of states between } k \text{ and } k + \mathrm{dk}) = g(k)\mathrm{dk}$$

and

$$g(k)\mathrm{dk} = \frac{\mathrm{V}}{(2\pi)^3} 4\pi \mathrm{k}^2 \mathrm{dk}$$

Statistical mechanics of a gas at “high” (i.e. not extremely low) temperature

1. Obtain partition function for the translational motion of particles in a box
2. Hence obtain energy, heat capacity, **and entropy**
3. **The idea of k-space and density of states**
4. → Bring in rotation and vibration

So far, we have treated translational motion.



For MONATOMIC gas that's the whole story

(except there can be fine structure in the internal electronic energy levels … see notes)

For DIATOMIC gas we have also rotation and vibration.



For POLYATOMIC gas there is even more rotation and vibration.



Heat capacity of a typical diatomic gas as a function of temperature.

Most gases have rotation but not vibration excited at room temperature.

Total heat capacity when rotation is fully excited

$$
\begin{aligned}
C_V &= C_V^{(t)} + C_V^{(r)} + C_V^{(v)} \\
&= \frac{3}{2}Nk_B + \begin{cases} Nk_B & (\text{linear molecule}) \\ \frac{3}{2}Nk_B & (\text{non}-\text{linear molecule}) \end{cases} + Nk_B \sum_{i=1}^{q_v} \frac{\left(T_{v,i}/(2T)\right)^2}{\sinh^2\left(T_{v,i}/(2T)\right)}
\end{aligned}
$$

Translational

rotational

Vibrational
(simple harmonic motion
for each normal mode)

H_2O



CO_2



Fig. E.1 Normal modes of H_2O and CO_2 molecules. The arrows indicate the motion during one half of any given oscillation. The second bending mode for CO_2 is in the direction perpendicular to the plane of the diagram.